



**LUND**  
UNIVERSITY



# Non-perturbative aspects of generators for eA

Leif Lönnblad

Department of Astronomy  
and Theoretical Physics  
Lund University

MC4EIC, 2021-11-18

# Outline

- ▶ DIS and the dipole picture
- ▶ Glauber calculations
- ▶ Collective effects
- ▶ Plans for PYTHIA8



# Why General Purpose Event Generators

(only three GPEGs: Herwig7, Pythia8, Sherpa)

We need event generators to model our data, but also to model the theory.

It's not enough to tune for one analysis/experiment, we need to tune to everything.

- ▶  $e^+e^- \Rightarrow$  Hadronisation and FSR
- ▶  $ep \Rightarrow$  ISR and remnant jets.
- ▶  $pp \Rightarrow$  UE and MPI
- ▶  $pA \Rightarrow$  small dense systems, flow
- ▶  $AA \Rightarrow$  large dense systems, jet quenching
- ▶  $eA \Rightarrow ?$

We need to understand what exactly we are measuring.



Are the *general purpose* event generators ready for the EIC?

Can they handle

- ▶  $eA$ ?
- ▶ DIS?
- ▶ Photo-production?
- ▶ Nuclei in general?
- ▶ Saturation? Polarisation? Lower energy? Diffraction? ...



Are the *general purpose* event generators ready for the EIC?

Can they handle

- ▶  $eA$ ?
- ▶ DIS?
- ▶ Photo-production?
- ▶ Nuclei in general?
- ▶ Saturation? Polarisation? Lower energy? Diffraction? ...



Are the *general purpose* event generators ready for the EIC?

Can they handle

- ▶  $eA$ ?
- ▶ DIS?
- ▶ Photo-production?
- ▶ Nuclei in general?
- ▶ Saturation? Polarisation? Lower energy? Diffraction? ...



Are the *general purpose* event generators ready for the EIC?

Can they handle

- ▶  $eA$ ?
- ▶ DIS?
- ▶ Photo-production?
- ▶ Nuclei in general?
- ▶ Saturation? Polarisation? Lower energy? Diffraction? ...



Are the *general purpose* event generators ready for the EIC?

Can they handle

- ▶  $eA$ ?
- ▶ DIS?
- ▶ Photo-production?
- ▶ Nuclei in general?
- ▶ Saturation? Polarisation? Lower energy? Diffraction? ...



Are the *general purpose* event generators ready for the EIC?

Can they handle

- ▶  $eA$ ?
- ▶ DIS?
- ▶ Photo-production?
- ▶ Nuclei in general?
- ▶ Saturation? Polarisation? Lower energy? Diffraction? ...



Are the *general purpose* event generators ready for the EIC?


Can they handle

- ▶  $eA$ ?
- ▶ DIS?
- ▶ Photo-production?
- ▶ Nuclei in general?
- ▶ Saturation? Polarisation? Lower energy? Diffraction? ...

Todo: EVERYTHING




# The standard DIS picture



- ▶ The Primary (hardest) vertex.
- ▶ Standard showers orders in  $k \perp$
- ▶ And allows the scattered electron to take recoil.  $x, Q^2$  not preserved in shower.
- ▶ No natural transition to photo-production for small  $Q^2$ .
- ▶ If we can do ep we can do eA




# The standard DIS picture



- ▶ The Primary (hardest) vertex.
- ▶ Standard showers orders in  $k \perp$
- ▶ And allows the scattered electron to take recoil.  $x, Q^2$  not preserved in shower.
- ▶ No natural transition to photo-production for small  $Q^2$ .
- ▶ If we can do ep we can do eA




# The standard DIS picture



- ▶ The Primary (hardest) vertex.
- ▶ Standard showers orders in  $k_{\perp}$
- ▶ And allows the scattered electron to take recoil.  $x, Q^2$  not preserved in shower.
- ▶ No natural transition to photo-production for small  $Q^2$ .
- ▶ If we can do ep we can do eA




# The standard DIS picture



- ▶ The Primary (hardest) vertex.
- ▶ Standard showers orders in  $k \perp$
- ▶ And allows the scattered electron to take recoil.  $x, Q^2$  not preserved in shower.
- ▶ No natural transition to photo-production for small  $Q^2$ .
- ▶ If we can do ep we can do eA




# The standard DIS picture



- ▶ The Primary (hardest) vertex.
- ▶ Standard showers orders in  $k \perp$
- ▶ And allows the scattered electron to take recoil.  $x, Q^2$  not preserved in shower.
- ▶ No natural transition to photo-production for small  $Q^2$ .
- ▶ If we can do ep we can do eA




# The standard DIS picture



- ▶ The Primary (hardest) vertex.
- ▶ Standard showers orders in  $k \perp$
- ▶ And allows the scattered electron to take recoil.  $x, Q^2$  not preserved in shower.
- ▶ No natural transition to photo-production for small  $Q^2$ .
- ▶ If we can do ep we can do eA




# The standard DIS picture



- ▶ The Primary (hardest) vertex.
- ▶ Standard showers orders in  $k \perp$
- ▶ And allows the scattered electron to take recoil.  $x, Q^2$  not preserved in shower.
- ▶ No natural transition to photo-production for small  $Q^2$ .
- ▶ If we can do ep we can do eA




## The standard DIS picture



- ▶ The Primary (hardest) vertex.
- ▶ Standard showers orders in  $k \perp$
- ▶ And allows the scattered electron to take recoil.  $x, Q^2$  not preserved in shower.
- ▶ No natural transition to photo-production for small  $Q^2$ .
- ▶ If we can do ep we can do eA?




# The dipole picture



- ▶ In the frame where the target is at rest the photon is emitted long before the interaction.
- ▶ Also the photon splits up long before the interaction
- ▶ ... and radiates
- ▶ ... before hitting the target.
- ▶ Hitting another nucleon comes at little extra cost.




# The dipole picture



- ▶ In the frame where the target is at rest the photon is emitted long before the interaction.
- ▶ Also the photon splits up long before the interaction
- ▶ ... and radiates
- ▶ ... before hitting the target.
- ▶ Hitting another nucleon comes at little extra cost.




# The dipole picture



- ▶ In the frame where the target is at rest the photon is emitted long before the interaction.
- ▶ Also the photon splits up long before the interaction
- ▶ ... and radiates
- ▶ ... before hitting the target.
- ▶ Hitting another nucleon comes at little extra cost.




# The dipole picture



- In the frame where the target is at rest the photon is emitted long before the interaction.
- Also the photon splits up long before the interaction
- ... and radiates
- ... before hitting the target.
- Hitting another nucleon comes at little extra cost.




# The dipole picture



- In the frame where the target is at rest the photon is emitted long before the interaction.
- Also the photon splits up long before the interaction
- ... and radiates
- ... before hitting the target.
- Hitting another nucleon comes at little extra cost.



# The dipole picture



- In the frame where the target is at rest the photon is emitted long before the interaction.
- Also the photon splits up long before the interaction
- ... and radiates
- ... before hitting the target.
- Hitting another nucleon comes at little extra cost.



# Glauber Calculations

To get a reasonable handle on hadronic final states in eA we need to add together several eN collisions, and for that we need to do some kind of Glauber calculation to obtain  $N_{\text{part}}$  (or  $N_{\text{wounded}}$ )

Also, if we want to do  $R_{AA}$ -like measurements, we need to understand how many nucleons we hit.

The Glauber calculation needs to use the dipole-nucleon cross section.



# Glauber Calculations

To get a reasonable handle on hadronic final states in eA we need to add together several eN collisions, and for that we need to do some kind of Glauber calculation to obtain  $N_{\text{part}}$  (or  $N_{\text{wounded}}$ )


Also, if we want to do  $R_{AA}$ -like measurements, we need to understand how many nucleons we hit.


The Glauber calculation needs to use the dipole-nucleon cross section.

The lesson from pA collisions at the LHC is that we need to worry about fluctuations.




# Number of wounded nucleons





For low  $Q^2$  eA we expect Vector Meson Dominance and the situation looks like pA.






For low  $Q^2$  eA we expect Vector Meson Dominance and the situation looks like pA.

For high  $Q^2$  we expect even more fluctuations. And we need to generate DIS in terms of dipole–nucleon scattering.





For low  $Q^2$  eA we expect Vector Meson Dominance and the situation looks like pA.

For high  $Q^2$  we expect even more fluctuations. And we need to generate DIS in terms of dipole–nucleon scattering.

More on this in Christian Bierlich's talk



# Hadronisation

We believe that we know how to model the non-perturbative hadronisation process. But there are caveats.

- ▶  $\sqrt{s_{eN}} = 100$  GeV, sounds like LEP energies  
But hadronic system has lower mass and a lots of energy goes into the nucleon remnant (largely ignored by GPEGs).
- ▶ How well does string fragmentation describe a small number of hadrons? Cluster fragmentation?
- ▶ How well do we model intrinsic  $k_{\perp}$ ?



# Hadronisation

We believe that we know how to model the non-perturbative hadronisation process. But there are caveats.

- ▶  $\sqrt{S_{eN}} = 100$  GeV, sounds like LEP energies  
But hadronic system has lower mass and a lots of energy goes into the nucleon remnant (largely ignored by GPEGs).
- ▶ How well does string fragmentation describe a small number of hadrons? Cluster fragmentation?
- ▶ How well do we model intrinsic  $k_{\perp}$ ?



# Hadronisation

We believe that we know how to model the non-perturbative hadronisation process. But there are caveats.

- ▶  $\sqrt{s_{eN}} = 100$  GeV, sounds like LEP energies  
But hadronic system has lower mass and a lots of energy goes into the nucleon remnant (largely ignored by GPEGs).
- ▶ How well does string fragmentation describe a small number of hadrons? Cluster fragmentation?
- ▶ How well do we model intrinsic  $k_{\perp}$ ?



# Hadronisation

We believe that we know how to model the non-perturbative hadronisation process. But there are caveats.

- ▶  $\sqrt{s_{eN}} = 100$  GeV, sounds like LEP energies  
But hadronic system has lower mass and a lots of energy goes into the nucleon remnant (largely ignored by GPEGs).
- ▶ How well does string fragmentation describe a small number of hadrons? Cluster fragmentation?
- ▶ How well do we model intrinsic  $k_{\perp}$ ?



# Collective Effects

Can we just stack dipole–nucleon events together, or do we need to worry about collective effects?



# Collective Effects

Can we just stack dipole–nucleon events together, or do we need to worry about collective effects?

Do we need to model a droplet of Quark–Gluon Plasma?



# Collective Effects

Can we just stack dipole–nucleon events together, or do we need to worry about collective effects?

Do we need to model a droplet of Quark–Gluon Plasma?

Since there are collective effects even in pp, we need to be able to model them at the EIC!



# Collective effects in PYTHIA8 (Gleipnir)

- ▶ Colour reconnections (swing)
- ▶ String repulsion (shoving)
- ▶ Increased string tension (ropes)
- ▶ Hadron rescattering



# Collective effects in PYTHIA8 (Gleipnir)

- ▶ Colour reconnections (swing)  
Affects jet shapes and multiplicity
- ▶ String repulsion (shoving)
- ▶ Increased string tension (ropes)
- ▶ Hadron rescattering



# Collective effects in PYTHIA8 (Gleipnir)

- ▶ Colour reconnections (swing)  
Affects jet shapes and multiplicity
- ▶ String repulsion (shoving)  
Gives flow
- ▶ Increased string tension (ropes)
- ▶ Hadron rescattering



# Collective effects in PYTHIA8 (Gleipnir)

- ▶ Colour reconnections (swing)  
Affects jet shapes and multiplicity
- ▶ String repulsion (shoving)  
Gives flow
- ▶ Increased string tension (ropes)  
Gives strangeness enhancement
- ▶ Hadron rescattering





# Collective effects in PYTHIA8 (Gleipnir)






- ▶ Colour reconnections (swing)  
Affects jet shapes and multiplicity
- ▶ String repulsion (shoving)  
Gives flow
- ▶ Increased string tension (ropes)  
Gives strangeness enhancement
- ▶ Hadron rescattering  
More flow




# Shoving (pp)



# Shoving (pPb)



# Ropes



Nature Phys. 13 (2017) 535-539



# EIC plans for PYTHIA8

- ▶ eA (Dipole–nucleon scatterings) in Angantyr [talk this afternoon]
- ▶ Swing
- ▶ Shoving
- ▶ Ropes
- ▶ Hadronic rescattering
- ▶ Nuclear PDFs
- ▶ Photon PDFs
- ▶ Polarised string fragmentation (w. Albi Kerbizi)
- ▶ TMD-based shower (w. Mees van Kampen)



# EIC plans for PYTHIA8

- ▶ eA (Dipole–nucleon scatterings) in Angantyr [talk this afternoon]
- ▶ Swing
- ▶ Shoving
- ▶ Ropes
- ▶ Hadronic rescattering
- ▶ Nuclear PDFs
- ▶ Photon PDFs
- ▶ Polarised string fragmentation (w. Albi Kerbizi)
- ▶ TMD-based shower (w. Mees van Kampen)



# EIC plans for PYTHIA8

- ▶ eA (Dipole–nucleon scatterings) in Angantyr [talk this afternoon]
- ▶ Swing
- ▶ Shoving
- ▶ Ropes
- ▶ Hadronic rescattering
- ▶ Nuclear PDFs
- ▶ Photon PDFs
- ▶ Polarised string fragmentation (w. Albi Kerbizi)
- ▶ TMD-based shower (w. Mees van Kampen)



# EIC plans for PYTHIA8

- ▶ eA (Dipole–nucleon scatterings) in Angantyr [talk this afternoon]
- ▶ Swing
- ▶ Shoving
- ▶ Ropes
- ▶ Hadronic rescattering
- ▶ Nuclear PDFs
- ▶ Photon PDFs
- ▶ Polarised string fragmentation (w. Albi Kerbizi)
- ▶ TMD-based shower (w. Mees van Kampen)



# EIC plans for PYTHIA8

- ▶ eA (Dipole–nucleon scatterings) in Angantyr [talk this afternoon]
- ▶ Swing
- ▶ Shoving
- ▶ Ropes
- ▶ Hadronic rescattering
- ▶ Nuclear PDFs
- ▶ Photon PDFs
- ▶ Polarised string fragmentation (w. Albi Kerbizi)
- ▶ TMD-based shower (w. Mees van Kampen)



# The Conclusion

General Purpose Event Generators are not ready for EIC.



# The Conclusion

General Purpose Event Generators are not ready for EIC. Yet.



# The Conclusion

General Purpose Event Generators are not ready for EIC. Yet.  
But there are at least PYTHIA8 authors that are willing to have a go at it.



# Thanks!



European Research Council  
Established by the European Commission

